

Lecture 43: Consistency of the IV estimator; an IV calculation

Econometrics ■ Unit III: Autocorrelation and instrumental variables

Kapil Kumar

Department of Statistics
Chaudhary Charan Singh University, Meerut

B.Sc. (Hons) Statistics, Semester V
Academic Session 2026–2027

Previous lecture

Lecture 42: Instruments; the IV estimator in the simple regression.

Today

- The IV estimator in the simple regression
 - Consistency

Substituting $y_i - \bar{y} = \beta_1(x_i - \bar{x}) + (\varepsilon_i - \bar{\varepsilon})$,

$$\hat{\beta}_{1,IV} = \beta_1 + \frac{\frac{1}{n} \sum_i (z_i - \bar{z}) \varepsilon_i}{\frac{1}{n} \sum_i (z_i - \bar{z})(x_i - \bar{x})}.$$

Taking probability limits,

$$\text{plim } \hat{\beta}_{1,IV} = \beta_1 + \frac{\text{Cov}(z, \varepsilon)}{\text{Cov}(z, x)} = \beta_1 + \frac{0}{\text{Cov}(z, x)} = \beta_1.$$

- So the IV estimator is consistent.
- It is generally *not* unbiased in finite samples.

Example 6.2 (An IV calculation)

i	z_i	x_i	y_i
1	-2	1	1
2	-1	2	3
3	0	2	2
4	1	4	5
5	2	6	9

Means. $\bar{z} = 0$, $\bar{x} = 15/5 = 3$, $\bar{y} = 20/5 = 4$.

Cross products. Since $\bar{z} = 0$,

$$\begin{aligned}\sum z_i(x_i - \bar{x}) &= \sum z_i x_i \\ &= -2 - 2 + 0 + 4 + 12 = 12,\end{aligned}$$

$$\begin{aligned}\sum z_i(y_i - \bar{y}) &= \sum z_i y_i \\ &= -2 - 3 + 0 + 5 + 18 = 18.\end{aligned}$$

Estimates. $\hat{\beta}_{1,IV} = 18/12 = 1.5$ and $\hat{\beta}_{0,IV} = 4 - 1.5(3) = -0.5$.

Example 6.2 (cont.)

Comparison with OLS.

$$\sum (x_i - \bar{x})^2 = 4 + 1 + 1 + 1 + 9 = 16,$$

$$\sum (x_i - \bar{x})(y_i - \bar{y}) = 6 + 1 + 2 + 1 + 15 = 25,$$

so $b_1 = 25/16 = 1.5625$ and $b_0 = 4 - 1.5625(3) = -0.6875$.

Residual check. The IV residuals $y_i + 0.5 - 1.5x_i$ are

$$(0, 0.5, -0.5, -0.5, 0.5).$$

They sum to zero and $\sum z_i e_i = 0 - 0.5 + 0 - 0.5 + 1 = 0$, as required by the IV equations.

Practice

- Exercise 6.3 (Exercises and Solutions booklet).

Next lecture

Lecture 44: The IV estimator in matrix form: consistency and variance.