

Lecture 58: LIML: the least variance ratio and its minimisation

Econometrics ■ Unit IV: Simultaneous equations models

Kapil Kumar

Department of Statistics
Chaudhary Charan Singh University, Meerut

B.Sc. (Hons) Statistics, Semester V
Academic Session 2026–2027

Previous lecture

Lecture 57: 2SLS standard errors; an over-identified example; 2SLS, IV and ILS under exact identification.

Today

- Limited information maximum likelihood
 - The idea: the least variance ratio
 - Minimising the variance ratio

- LIML is due to Anderson and Rubin (1949).
- It estimates one structural equation using only the information about that equation (its included and excluded variables), not the full specification of the other equations; hence “limited information”.
- Under normal disturbances, it is the maximum likelihood estimator based on this limited information.
- We derive it by the equivalent *least variance ratio* principle, which needs only calculus.

Let $H = [y_1 \ Y_1]$ (the $m + 1$ endogenous variables in the equation) and $\mathbf{a} = (1, -\beta')'$. For a trial value of β , define

$$y^* = y_1 - Y_1\beta = H\mathbf{a}.$$

- If β is the true value, then $y^* = X_1\gamma + u$: y^* depends only on the *included* exogenous variables.
- The excluded variables X_2 should then add nothing to explaining y^* .

Two regressions of y^*

- Regress y^* on X_1 only. Its residual sum of squares is (Preliminaries, $e'e = y'My$)

$$\text{RSS}_1 = y^{*'}M_1y^* = \mathbf{a}'H'M_1H\mathbf{a} = \mathbf{a}'W_1\mathbf{a},$$

where $M_1 = I - X_1(X_1'X_1)^{-1}X_1'$ and $W_1 = H'M_1H$.

- Regress y^* on all exogenous variables X . Its residual sum of squares is

$$\text{RSS} = \mathbf{a}'W\mathbf{a},$$

where $M = I - X(X'X)^{-1}X'$ and $W = H'MH$.

Since X contains X_1 , adding regressors cannot increase the residual sum of squares, so $\text{RSS}_1 \geq \text{RSS}$. Define the **variance ratio**

$$\kappa(\mathbf{a}) = \frac{\mathbf{a}'W_1\mathbf{a}}{\mathbf{a}'W\mathbf{a}} \geq 1.$$

If the excluded variables have no explanatory power for y^* , κ is close to 1.

LIML chooses β to make κ as small as possible.

Step 1 (differentiate). W_1 and W are symmetric, so by Rule P2 and the quotient rule,

$$\frac{\partial \kappa}{\partial \mathbf{a}} = \frac{2W_1\mathbf{a}(\mathbf{a}'W\mathbf{a}) - (\mathbf{a}'W_1\mathbf{a})2W\mathbf{a}}{(\mathbf{a}'W\mathbf{a})^2}.$$

Step 2 (equate to zero). Dividing by $2/(\mathbf{a}'W\mathbf{a})$,

$$W_1\mathbf{a} - \frac{\mathbf{a}'W_1\mathbf{a}}{\mathbf{a}'W\mathbf{a}}W\mathbf{a} = 0 \implies (W_1 - \kappa W)\mathbf{a} = 0.$$

Step 3 (determinantal equation). A nonzero α satisfies this only if

$$|W_1 - \kappa W| = 0.$$

This is a polynomial equation of degree $m + 1$ in κ . At each root, κ equals the value of the ratio at the corresponding α .

Step 4 (choose the smallest root). Since we want the minimum of the ratio, take the **smallest root** $\hat{\kappa}$.

Step 5 (coefficients). Solve $(W_1 - \hat{\kappa}W)\mathbf{a} = 0$, scale the solution so that its first element is 1, and read off $\hat{\beta}_{LIML}$ as minus the remaining elements. Then estimate γ by regressing $y_1 - Y_1\hat{\beta}_{LIML}$ on X_1 :

$$\hat{\gamma}_{LIML} = (X_1'X_1)^{-1}X_1'(y_1 - Y_1\hat{\beta}_{LIML}).$$

The matrices W_1 and W are only $(m+1) \times (m+1)$, whatever the sample size.

Practice

Re-work the derivations and examples of this lecture without looking at the slides.

Next lecture

Lecture 59: LIML under exact identification; the k -class; a LIML calculation.