

Lecture 3: Randomised decision rules; convexity of the risk set

Statistical Inference–II ■ Unit I: Decision Theory

Kapil Kumar

Department of Statistics
Chaudhary Charan Singh University, Meerut

M.Sc. Statistics, Semester III
Academic Session 2026–2027

Previous lecture

Lecture 2: Non-randomised decision rules; the risk function; risk of an estimator and of a test.

Today

- Decision Rules and the Risk Function
 - Randomised Decision Rules

In some problems it is advantageous — and in some it is unavoidable — to allow the action to be chosen by an auxiliary random mechanism after $\mathbf{x}$ has been observed. This is familiar from the theory of testing, where randomised tests are needed to attain an exact size α in discrete problems.

Definition 3.1 (Randomised decision rule)

A *randomised decision rule* δ^* is a function which assigns to each $\mathbf{x} \in \mathcal{X}$ a probability distribution $\delta^*(\cdot \mid \mathbf{x})$ on the action space $\mathcal{A}$. When $\mathbf{x}$ is observed, the action actually taken is drawn from $\delta^*(\cdot \mid \mathbf{x})$ by an auxiliary randomisation whose law does not depend on θ .

When $\mathcal{A} = \{a_1, a_2, \dots, a_k\}$ is finite, δ^* is described by k functions $p_1(\mathbf{x}), \dots, p_k(\mathbf{x})$ with

$$p_j(\mathbf{x}) \geq 0 \quad \text{and} \quad \sum_j p_j(\mathbf{x}) = 1,$$

where $p_j(\mathbf{x})$ is the probability of taking action a_j when $\mathbf{x}$ is observed.

- A non-randomised rule is the special case in which, for each $\mathbf{x}$, some $p_j(\mathbf{x}) = 1$.

For the two-action problem it is customary to write a randomised rule as a single *test function*

$$\phi(\mathbf{x}) = \mathbb{P}(\text{take action } a_1 \mid \mathbf{X} = \mathbf{x}) \in [0, 1].$$

Definition 3.2 (Risk of a randomised rule)

The loss incurred by a randomised rule at $(\theta, \mathbf{x})$ is itself an expected value over the randomisation:

$$L(\theta, \delta^*(\cdot | \mathbf{x})) = \int_{\mathcal{A}} L(\theta, a) \delta^*(da | \mathbf{x}).$$

The risk function is therefore

$$R(\theta, \delta^*) = \int_{\mathcal{X}} \left[\int_{\mathcal{A}} L(\theta, a) \delta^*(da | \mathbf{x}) \right] f(\mathbf{x} | \theta) d\mathbf{x}.$$

In the finite-action case this reduces to

$$R(\theta, \delta^*) = \mathbb{E}_{\theta} \left[\sum_{j=1}^k L(\theta, a_j) p_j(\mathbf{X}) \right].$$

Proposition 3.3 (Randomisation convexifies the risk set)

Let $\delta_1, \delta_2 \in \mathcal{D}$ and let δ^ be the randomised rule which uses δ_1 with probability λ and δ_2 with probability $1 - \lambda$, $0 \leq \lambda \leq 1$, the choice being made independently of $\mathbf{X}$. Then*

$$R(\theta, \delta^*) = \lambda R(\theta, \delta_1) + (1 - \lambda)R(\theta, \delta_2), \quad \forall \theta \in \Theta.$$

Proof.

Let U be a Bernoulli(λ) random variable independent of $\mathbf{X}$, and let δ^* use δ_1 if $U = 1$ and δ_2 if $U = 0$. Conditioning on U ,

$$\begin{aligned} R(\theta, \delta^*) &= \mathbb{E}_\theta[L(\theta, \delta^*(\mathbf{X}, U))] \\ &= \mathbb{E}\{\mathbb{E}_\theta[L(\theta, \delta^*(\mathbf{X}, U)) \mid U]\} \\ &= \mathbb{P}(U = 1) \mathbb{E}_\theta[L(\theta, \delta_1(\mathbf{X}))] + \mathbb{P}(U = 0) \mathbb{E}_\theta[L(\theta, \delta_2(\mathbf{X}))] \\ &= \lambda R(\theta, \delta_1) + (1 - \lambda) R(\theta, \delta_2). \end{aligned}$$



The proposition says that the set of attainable risk functions becomes *convex* once randomised rules are admitted. This has two important consequences.

- (i) **Better minimax rules.** Mixing two rules can lower the maximum risk: if δ_1 is good where δ_2 is bad and vice versa, a suitable mixture balances the two.
- Hence the smallest attainable maximum risk over randomised rules can be strictly smaller than over non-randomised rules.
 - This is illustrated in “Randomised Minimax Rules in Finite Problems” and in end-of-chapter Exercise 1.20, where the minimax rule is necessarily randomised.

(The game-theoretic background, the *minimax theorem*, is outlined in Lecture Notes, Appendix A.)

- (ii) **Randomisation never helps under a convex loss.** If $\mathcal{A}$ is a convex subset of $\mathbb{R}$ and $L(\theta, \cdot)$ is convex for each θ , then by Jensen's inequality the non-randomised rule

$$\bar{\delta}(\mathbf{x}) = \int a \delta^*(da \mid \mathbf{x})$$

satisfies

$$R(\theta, \bar{\delta}) \leq R(\theta, \delta^*) \quad \text{for all } \theta.$$

Hence in estimation under squared error loss one may restrict attention to non-randomised rules without loss. This is the content of the Rao–Blackwell type argument and explains why randomisation is prominent in testing but invisible in estimation.

Practice (Lecture Notes, Chapter 1)

- Short-answer questions (Section 1.12): Q2, Q15.

Next lecture

Lecture 4: Uniformly best rules; admissibility; linear estimators of a normal mean.