

Lecture 40: Summary and revision

Statistical Inference–II ■ Units II–III: Sequential Analysis

Kapil Kumar

Department of Statistics
Chaudhary Charan Singh University, Meerut

M.Sc. Statistics, Semester III
Academic Session 2026–2027

This lecture

Previous lecture

Lecture 39: Solved numerical examples: exponential and gamma.

Today

- Summary and Formula Sheet
- Short-Answer Questions

Key ideas of Units II–III

1. In a sequential test the sample size N is a random stopping time. The SPRT continues while the log-likelihood ratio W_m lies between $\log B$ and $\log A$.
2. W_m is a random walk; under H_0 it drifts down, under H_1 it drifts up. The SPRT terminates with probability one, and all moments of N are finite.
3. Wald's boundaries $A \approx (1 - \beta)/\alpha$, $B \approx \beta/(1 - \alpha)$ are safe: the attained error probabilities can exceed the nominal ones only slightly, and their sum cannot exceed $\alpha + \beta$.

Key ideas of Units II–III (cont.)

4. Wald's fundamental identity gives the O.C. function; Wald's equation gives the A.S.N. function; Wald's second identity handles the zero-drift point. The only approximation is neglect of the overshoot.
5. For one-parameter exponential families the SPRT reduces to two parallel straight lines for a running total S_m .
6. The SPRT typically needs far fewer observations on average than the fixed-sample test with the same error probabilities.

Quantity	Formula
Increments and walk	$Z_i = \log\{f(X_i; \theta_1)/f(X_i; \theta_0)\},$ $W_m = \sum_{i=1}^m Z_i = \log \Lambda_m$
SPRT	accept H_0 if $W_m \leq \log B$; reject if $W_m \geq \log A$; continue otherwise
Boundaries	$A \approx \frac{1 - \beta}{\alpha}, \quad B \approx \frac{\beta}{1 - \alpha}$
Wald's bounds	$\alpha' \leq \frac{\alpha}{1 - \beta}, \quad \beta' \leq \frac{\beta}{1 - \alpha}, \quad \alpha' + \beta' \leq \alpha + \beta$
Termination	$\mathbb{P}_\theta(N > n) \leq \rho^{\lfloor n/k \rfloor}, \quad 0 < \rho < 1; \quad \mathbb{E}_\theta(N^r) < \infty$

Quantity	Formula
Fundamental identity	$\mathbb{E}[e^{tW_N} \phi(t)^{-N}] = 1, \quad \phi(t) = \mathbb{E}(e^{tZ_1}) \text{ (} N \text{ bounded; or SPRT with } \phi(t) \geq 1)$
Wald's equation	$\mathbb{E}(W_N) = \mathbb{E}(N) \mathbb{E}(Z_1)$
Second identity	$\mathbb{E}(W_N^2) = \mathbb{E}(N) \mathbb{E}(Z_1^2) \text{ when } \mathbb{E}(Z_1) = 0$
O.C. root	$\phi_\theta(h) = 1, h \neq 0; \quad h(\theta_0) = 1, h(\theta_1) = -1$
O.C. function	$Q(\theta) \approx \frac{A^h - 1}{A^h - B^h}; \quad \text{at } h = 0: Q \approx \frac{\log A}{\log A - \log B}$

Quantity	Formula
A.S.N. function	$\mathbb{E}_{\theta}(N) \approx \frac{Q \log B + (1 - Q) \log A}{\mu_Z(\theta)},$ $\mu_Z(\theta) = \mathbb{E}_{\theta}(Z_1) \neq 0$
Zero drift	$\mathbb{E}_{\theta^*}(N) \approx -\frac{(\log A)(\log B)}{\mathbb{E}_{\theta^*}(Z_1^2)}$
Bernoulli lines	<p>continue while $-\ell_0 + s m < T_m < \ell_1 + s m$,</p> $s = \frac{g_2}{g_1 + g_2}, \ell_0 = \frac{-\log B}{g_1 + g_2}, \ell_1 = \frac{\log A}{g_1 + g_2}$
Other models	see the summary tables of common SPRTs (section on distribution-specific SPRTs)

Short-answer questions (Q1–Q7)

- Q1. What is a sequential test? How does it differ from a fixed-sample test?
- Q2. Define a stopping time. Why must the decision to stop at stage m depend only on $X_1, \dots, X_m$?
- Q3. Define Wald's SPRT for testing a simple hypothesis against a simple alternative.
- Q4. Write Wald's approximations to the boundaries A and B in terms of α and β .
- Q5. State Wald's bounds on the attained error probabilities.
- Q6. Why are the attained error probabilities of an SPRT usually smaller than the nominal ones?
- Q7. State the theorem on termination of the SPRT with probability one.

Short-answer questions (Q8–Q14)

- Q8. State Wald's fundamental identity and Wald's equation.
- Q9. Define the O.C. and A.S.N. functions of an SPRT.
- Q10. What are the values of $h(\theta)$ at θ_0 and θ_1 , and what are $Q(\theta_0)$ and $Q(\theta_1)$?
- Q11. Why can the usual A.S.N. formula not be used at the zero-drift point? What replaces it?
- Q12. Show that the SPRT for a normal mean depends on the data only through $\sum X_i$.
- Q13. Why does the O.C. curve of the gamma SPRT not depend on the shape parameter?
- Q14. Interpret the sign of $\mathbb{E}_{\theta_0}(Z_1) = -K(f_0, f_1)$.

Practice (Lecture Notes, Chapter 2)

- Short-answer questions (Section 2.12): Q1–Q14 (all, for revision).

Next lecture

Lecture 41 begins Units IV–V: Nonparametric Inference.