

Lecture 47: Runs: the null distribution of the number of runs

Statistical Inference–II ■ Units IV–V: Nonparametric Inference

Kapil Kumar

Department of Statistics
Chaudhary Charan Singh University, Meerut

M.Sc. Statistics, Semester III
Academic Session 2026–2027

Previous lecture

Lecture 46: One-sided bounds; duality with the sign test; principles of nonparametric testing.

Today

- Tests for Randomness Based on Runs
 - Conditional Distribution of the Number of Runs

A *run* is a maximal uninterrupted sequence of identical symbols. For example,

$$A A B B A B B$$

contains four runs: AA , BB , A and BB .

Runs tests detect clustering or excessive alternation.

- Too few runs indicate clustering or persistence;
- too many runs indicate rapid alternation.

Suppose a sequence contains n_1 symbols of type A , n_2 symbols of type B , and $n = n_1 + n_2$. Conditional on (n_1, n_2) , all $\binom{n}{n_1}$ arrangements are equally likely under randomness.

Lemma 47.1 (Compositions)

The number of ways of dividing r identical symbols, placed in a row, into k non-empty runs is $\binom{r-1}{k-1}$.

Proof.

A division into k runs is obtained by choosing $k - 1$ of the $r - 1$ gaps between adjacent symbols at which to cut, and every choice gives a different division. □

Theorem 47.2 (Null distribution of the number of runs)

Under randomness, conditionally on (n_1, n_2) ,

$$\mathbb{P}(R = 2k) = \frac{2 \binom{n_1-1}{k-1} \binom{n_2-1}{k-1}}{\binom{n}{n_1}},$$

$$\mathbb{P}(R = 2k + 1) = \frac{\binom{n_1-1}{k} \binom{n_2-1}{k-1} + \binom{n_1-1}{k-1} \binom{n_2-1}{k}}{\binom{n}{n_1}},$$

where a binomial coefficient with an invalid lower index is interpreted as zero.

Proof of Theorem: Null distribution of the number of runs

Proof.

Runs of A 's and B 's must alternate, and an arrangement is completely determined by the symbol it starts with and by the lengths of its successive runs.

Even case, $R = 2k$. There are k runs of each type. The sequence may start with A or with B (2 ways); the n_1 A 's can be split into k runs in $\binom{n_1-1}{k-1}$ ways and the n_2 B 's in $\binom{n_2-1}{k-1}$ ways (Compositions Lemma).

Odd case, $R = 2k + 1$. Either there are $k + 1$ runs of A and k runs of B (the sequence must then start and end with A), giving $\binom{n_1-1}{k} \binom{n_2-1}{k-1}$ arrangements, or k runs of A and $k + 1$ runs of B , giving $\binom{n_1-1}{k-1} \binom{n_2-1}{k}$ arrangements.

Dividing by the total number $\binom{n}{n_1}$ of equally likely arrangements gives the result. □

Practice (Lecture Notes, Chapter 3)

- Short-answer questions (Section 3.14): Q5.
- Exercises (Section 3.15): 3.8.

Next lecture

Lecture 48: Mean and variance of the number of runs; the one-sample runs test.