

Data Analysis using R

Dr. Kapil Kumar

Assistant Professor, Department of Statistics
Central University of Haryana, Mahendergarh, India
kapilstats@gmail.com

9 February 2019

One-day Workshop on Data Analysis Using R (Tratrix Club)
Department of Mathematics & Statistics, Manipal University Jaipur
Invited lecture

Plan of the session

- 1 Getting started with R and RStudio
- 2 Vectors and data objects
- 3 Graphics in base R
- 4 Categorical data
- 5 Comparing groups: t -tests and ANOVA
- 6 Correlation and regression

By the end you can install R and RStudio, create and summarise vectors, read a CSV file, draw basic plots, and run and interpret a chi-square test, t -tests, a one-way ANOVA and a simple linear regression.

- 1 Getting started
- 2 Vectors and data
- 3 Graphics
- 4 Categorical data
- 5 Comparing groups
- 6 Correlation and regression

Installing R and RStudio


- 1 Install R first, from CRAN:
<https://cran.r-project.org>
- 2 Then install the free RStudio Desktop:
<https://posit.co/download/rstudio-desktop/>
- 3 Open **RStudio** (not R) for daily work; it finds R automatically.

Both are free and run on Windows, macOS and Linux.

Engine and dashboard R is the engine that does the computing. RStudio is the dashboard around it: an editor, a console, a plot window and a file browser in one place.

Then Install extra packages once with `install.packages()`; load them in every session with `library()`.

Where R came from



The name R is named partly after the first names of its two authors (Ross and Robert) and partly as a play on the name S.

- Free and open-source software
- A full programming language: conditionals, loops, user-defined (even recursive) functions, input and output
- Excellent graphics for data exploration and publication
- The Comprehensive R Archive Network (CRAN) stores identical, up-to-date copies of R, its packages and documentation on servers worldwide
- On 7 February 2019 CRAN listed **13,689 packages**; 38 *Task Views* group them by topic

More than a statistics package

R is best thought of as an *environment* in which statistical techniques are implemented. Anyone can add new methods as a package.


Packages:

<https://cran.r-project.org/web/packages/>

Task Views:

<https://cran.r-project.org/web/views/>

The four panes of RStudio



- Write in the script; use the console only for quick checks
- Start comments with #
- Keep the script and its data in one folder (an RStudio *project*)
- *Import Dataset* reads CSV, Excel, SPSS, SAS and Stata files and shows the code it used

Packages used in this session

- Base R already contains everything for today: `read.csv()`, `table()`, `chisq.test()`, `t.test()`, `aov()`, `lm()`, `plot()`
- **arm**: `display()` prints a compact summary of a fitted regression
- Install a package once (internet needed); load it in each new session

```
# install once  
# install.packages("arm")  
library(arm)
```

Tip `library()` without arguments lists the packages installed on your computer.

- 1 Getting started
- 2 Vectors and data**
- 3 Graphics
- 4 Categorical data
- 5 Comparing groups
- 6 Correlation and regression

R as a calculator

Operator	Meaning
+ -	add, subtract
* /	multiply, divide
^	power
%%	integer division
%%	remainder (modulo)
()	group: evaluated first

As in school algebra, powers come before * and /, which come before + and -.

```
2 + 2; 5 - 3; 5 / 2; 5^2
```

```
## [1] 4
```

```
## [1] 2
```

```
## [1] 2.5
```

```
## [1] 25
```

```
3 + 2 * 5; (3 + 2) * 5 # * before +; brackets first
```

```
## [1] 13
```

```
## [1] 25
```

```
7 %/% 2; 7 %% 2 # integer part, remainder
```

```
## [1] 3
```

```
## [1] 1
```

A data vector and its summary statistics

```
x <- c(2, 4, 3, 5, 7, 9, 15) # c() combines  
length(x); sum(x); mean(x); median(x)
```

```
## [1] 7  
## [1] 45  
## [1] 6.429  
## [1] 5
```

```
var(x); sd(x); range(x)
```

```
## [1] 19.95  
## [1] 4.467  
## [1] 2 15
```

```
summary(x) # quantile(x) gives the quartiles
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.  
##      2.00   3.50   5.00   6.43   8.00   15.00
```

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$
$$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$$

- `<-` assigns: read it as “gets”
- `var()` and `sd()` use the divisor $n - 1$
- Mean 6.43 > median 5: the value 15 pulls the mean up (right skew)

Vector arithmetic works element by element

- A function applied to a vector acts on *every* element: no loop needed
- `log()` is the natural log; `log10()`, `log2()` or the base argument give others
- `gamma()` is the gamma function: $\Gamma(n) = (n - 1)!$
- `round(v, 2)` keeps two decimals; `sqrt()`, `exp()`, `sin()`, `cos()` work the same way

```
1 + x; 2 * x; x^2
```

```
## [1] 3 5 4 6 8 10 16
```

```
## [1] 4 8 6 10 14 18 30
```

```
## [1] 4 16 9 25 49 81 225
```

```
round(log(x), 2) # natural log
```

```
## [1] 0.69 1.39 1.10 1.61 1.95 2.20 2.71
```

```
round(log(x, base = 10), 2) # same as log10(x)
```

```
## [1] 0.30 0.60 0.48 0.70 0.85 0.95 1.18
```

```
gamma(x)
```

```
## [1] 1 6 2 24
```

```
## [5] 720 40320 87178291200
```

Sorting, ordering and extracting elements

```
sort(x); rev(sort(x)) # up, then down
```

```
## [1] 2 3 4 5 7 9 15
```

```
## [1] 15 9 7 5 4 3 2
```

```
order(x); x[order(x)] # positions; sorted
```

```
## [1] 1 3 2 4 5 6 7
```

```
## [1] 2 3 4 5 7 9 15
```

sort or order? `sort()` returns the values; `order()` returns their positions, which can be used to reorder other columns too.

```
x[2]; x[c(2, 4)] # pick elements
```

```
## [1] 4
```

```
## [1] 4 5
```

```
x[-2]; x[-c(1, 3)] # drop elements
```

```
## [1] 2 3 5 7 9 15
```

```
## [1] 4 5 7 9 15
```

```
x[x > 5]; which(x > 5) # by a condition
```

```
## [1] 7 9 15
```

```
## [1] 5 6 7
```

[] extracts; a minus sign drops; a condition keeps the TRUE ones.

Sequences and combining vectors

```
3:6; seq(from = 3, to = 10)

## [1] 3 4 5 6
## [1] 3 4 5 6 7 8 9 10

seq(from = 3, to = 10, by = 0.5)

## [1] 3.0 3.5 4.0 4.5 5.0 5.5 6.0 6.5
## [9] 7.0 7.5 8.0 8.5 9.0 9.5 10.0

seq(from = 3, to = 5, length.out = 5)

## [1] 3.0 3.5 4.0 4.5 5.0

rep(c(2, 3), times = c(5, 4)); rep(1:3, 3)

## [1] 2 2 2 2 2 3 3 3 3
## [1] 1 2 3 1 2 3 1 2 3
```

```
x1 <- c(1, 2, 3, 4, 5, 6, 7)
y1 <- c(2, 1, 4, 3, 2, 5, 6)
c(x1, y1)                                # one long vector

## [1] 1 2 3 4 5 6 7 2 1 4 3 2 5 6

rbind(x1, y1)                             # as rows

##      [,1] [,2] [,3] [,4] [,5] [,6] [,7]
## x1      1      2      3      4      5      6      7
## y1      2      1      4      3      2      5      6

dim(cbind(x1, y1))                        # as columns: 7 x 2

## [1] 7 2
```

Naming objects and naming a vector

- A name starts with a letter (or a dot not followed by a digit) and may contain letters, digits, . and _
- R is case sensitive: x and X are different objects
- Avoid names of existing functions such as c, t or mean
- The elements of a vector can carry names too

```
names(x) <- c("A", "B", "C", "D",  
              "E", "F", "G")
```

```
x; x["D"]
```

```
##  A  B  C  D  E  F  G  
##  2  4  3  5  7  9 15  
## D  
## 5
```

Data objects vectors, matrices, data frames, lists and factors.

Lists and factors

A **list** collects objects of different types and lengths.

```
data5 <- list(x = c(2, 5, 4),
              Name = c("X", "Y", "Z"),
              Median = 4)
str(data5)

## List of 3
## $ x      : num [1:3] 2 5 4
## $ Name   : chr [1:3] "X" "Y" "Z"
## $ Median: num 4

data5$Name; data5[["Median"]]

## [1] "X" "Y" "Z"
## [1] 4
```

A **factor** stores a categorical variable with its levels.

```
group <- gl(2, 5, labels = c("batchA",
                             "batchB"))
group

## [1] batchA batchA batchA batchA batchA
## [6] batchB batchB batchB batchB batchB
## Levels: batchA batchB

levels(group); table(group)

## [1] "batchA" "batchB"
## group
## batchA batchB
##      5      5
```

Getting help in R

```
help(mean)           # or ?mean
??correlation        # search all packages
help.search("correlation")
help(package = "MASS")
example(mean)        # run the examples
?swiss               # describe a data set
```

- `?` opens the help page of a function you know by name
- `??` searches every installed package for a word
- `data()` lists the data sets that come with R
- `str()` shows the structure of any object

```
str(sleep)  # a built-in data frame: 20 rows, 3 columns

## 'data.frame': 20 obs. of  3 variables:
## $ extra: num  0.7 -1.6 -0.2 -1.2 -0.1 3.4 3.7 0.8 0 2 ...
## $ group: Factor w/ 2 levels "1","2": 1 1 1 1 1 1 1 1 1 1 ...
## $ ID   : Factor w/ 10 levels "1","2","3","4",...: 1 2 3 4 5 6 7 8 9 10 ...
```

Importing and exporting a CSV file

- Enter data in Excel, one row per case and variable names in the first row; save as CSV (*comma delimited*)
- `read.csv()` reads it into a data frame
- `write.csv()` saves a data frame
- Use relative paths such as `data/...`; avoid hard-coded paths like `D:/...`

```
data1 <- read.csv("data/data1.csv")
data1

##      A  B C  D
## 1  10 12 1 14
## 2  25 24 2 36
## 3  36 36 5 69
## 4  14 85 9 58

write.csv(data1, "data/data1_copy.csv",
          row.names = FALSE)
copy1 <- read.csv("data/data1_copy.csv")
identical(data1, copy1) # same data back?

## [1] TRUE
```

`data1.csv`: four illustrative variables A–D, four rows.

apply(): summaries by row or by column

```
apply(data1, 1, mean)      # row means

## [1]  9.25 21.75 36.50 41.50

apply(data1, 2, mean)      # column means

##      A      B      C      D
## 21.25 39.25  4.25 44.25

apply(data1, 2, sd)

##      A      B      C      D
## 11.701 32.035  3.594 24.391

apply(data1, 2, median)

##      A      B      C      D
## 19.5 30.0  3.5 47.0
```

`apply(X, MARGIN, FUN)` `MARGIN`
= 1 works along rows, `MARGIN` = 2
along columns; `FUN` is any
function that takes a vector.

Shortcuts for the most common
cases: `rowMeans()`, `colMeans()`,
`rowSums()`, `colSums()`.

Writing your own function: CV and standard error

$$CV = 100 \frac{s}{\bar{x}} \%, \quad SE(\bar{x}) = \frac{s}{\sqrt{n}}$$

- CV compares spread across variables measured on different scales (positive data only)
- SE measures how precisely $\bar{x}$ estimates the population mean

```
cv <- function(v) 100 * sd(v) / mean(v)
se <- function(v) sd(v) / sqrt(length(v))
round(apply(data1, 2, cv), 1)
```

```
##      A      B      C      D
## 55.1 81.6 84.6 55.1
```

```
round(apply(data1, 2, se), 2)
```

```
##      A      B      C      D
## 5.85 16.02 1.80 12.20
```

Reading it C has the smallest SE (1.8) but the largest CV (84.6%): small values, yet large *relative* spread.

- 1 Getting started
- 2 Vectors and data
- 3 Graphics**
- 4 Categorical data
- 5 Comparing groups
- 6 Correlation and regression

High level

start a new plot

`plot()`, `hist()`
`boxplot()`, `barplot()`
`dotchart()`, `qqnorm()`
`pie()`, `curve()`

Low level

add to the current plot

`points()`, `lines()`
`abline()`, `text()`
`grid()`, `legend()`
`qqline()`, `title()`

`par(mfrow = c(r, c))` puts $r \times c$ plots on one page; `par(mar = ...)` sets the margins.

```
length(colors()) # named colours
```

```
## [1] 657
```

```
colors()[c(26, 254, 552)]
```

```
## [1] "blue" "green" "red"
```




```
# colours for today (hex codes)
```

```
col_main <- "#3333B3" # blue
```

```
col_alt <- "#B45309" # amber
```

Bar plots and dot charts of named scores

```
par(mfrow = c(1, 3), mar = c(4, 4, 2, 1))  
barplot(x, las = 1, col = col_main, main = "Scores of students")  
barplot(sort(x), las = 1, col = col_alt, main = "Sorted by score")  
dotchart(sort(x), pch = 16, col = col_main, main = "Dot chart")
```




`las = 1` keeps axis labels horizontal. A dot chart shows the same information with less ink and stays readable with many categories.

Building a plot layer by layer


```
x2 <- c(1, 2, 3, 4, 5, 6, 7)
y2 <- c(2, 1, 4, 3, 2, 5, 6)
par(mar = c(4, 4, 2, 1))
plot(x2, y2, type = "b", pch = 5, lty = 4,
     col = col_main, main = "My plot",
     xlab = "x axis", ylab = "y axis")
grid(col = "grey80")
text(2.5, 5, "this is my example plot")
abline(h = 2, v = 4, col = c(col_alt,
                              "darkgreen"), lty = c(1, 4), lwd = 2)
```

type: "p" points, "l" lines, "b" both; pch: symbol;
lty: line type; lwd: line width.



Is a sample normal? Histogram and normal Q-Q plot

```
set.seed(2019)
rn <- rnorm(200); rb <- rbinom(20, 100, 0.85) # N(0, 1) and Binomial(100, 0.85) samples
par(mfrow = c(1, 4), mar = c(4, 4, 2, 1))
hist(rn, col = col_main); qqnorm(rn); qqline(rn, col = col_alt, lwd = 2)
hist(rb, col = col_alt); qqnorm(rb); qqline(rb, col = col_main, lwd = 2)
```



Points close to the line suggest normality. The binomial counts are whole numbers, so tied values form horizontal steps; with only 20 values the histogram is rough.

- 1 Getting started
- 2 Vectors and data
- 3 Graphics
- 4 Categorical data**
- 5 Comparing groups
- 6 Correlation and regression

The sugar data

```
data4 <- read.table("data/data4.txt",
                    header = TRUE)

str(data4)

## 'data.frame': 20 obs. of 3 variables:
## $ Sugar : chr  "no" "no" "yes" "yes" ...
## $ Age   : chr  "old" "young" "old" "old" ...
## $ Gender: chr  "female" "female" "male" "male" ...

head(data4, 3)

##   Sugar   Age Gender
## 1    no   old female
## 2    no young female
## 3   yes   old  male
```

- Hypothetical data on 20 patients
- Sugar: diabetic (yes/no); Age: old/young; Gender
- `read.table()` reads text files whose columns are separated by spaces; `header = TRUE` takes names from the first row

Question Is having sugar associated with age group, or with gender?

Frequency tables

```
table(data4$Sugar)
```

```
##  
##  no  yes  
## 13   7
```

```
with(data4, table(Sugar, Gender))
```

```
##      Gender  
## Sugar female male  
##   no         6   7  
##   yes         5   2
```

`with(data4, ...)` lets us write Sugar instead of `data4$Sugar`.

```
with(data4, table(Sugar, Age))
```

```
##      Age  
## Sugar old young  
##   no    2    11  
##   yes    6     1
```

```
ftable(with(data4, table(Gender, Age, Sugar)))
```

```
##              Sugar no yes  
## Gender Age  
## female old          1   4  
##        young        5   1  
## male   old          1   2  
##        young        6   0
```

Row proportions tell the story


```
tab <- with(data4, table(Age, Sugar))
addmargins(tab) # with totals

##           Sugar
## Age      no yes Sum
##  old       2  6  8
##  young    11  1 12
##  Sum      13  7 20

round(prop.table(tab, 1), 2) # 1 = rows, 2 = cols

##           Sugar
## Age      no yes
##  old    0.25 0.75
##  young  0.92 0.08
```

```
p_yes <- prop.table(tab, 1)[, "yes"]
par(mar = c(4, 4, 1, 1))
barplot(p_yes, col = col_main, las = 1,
        ylim = c(0, 1),
        ylab = "Proportion with sugar")
```



75% of the old patients have sugar,
against 8% of the young ones.

Chi-square test of independence, and a small-sample check

H_0 : sugar status and age group are independent

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

$E = \text{row total} \times \text{column total} / n$; $df = (r - 1)(c - 1)$

Rule of thumb The χ^2 approximation needs expected counts $E \geq 5$. Two are smaller here (R warns that the approximation may be incorrect), so use Fisher's exact test.

```
ct <- chisq.test(tab)
round(ct$expected, 1)      # expected counts

##           Sugar
## Age         no yes
##   old      5.2 2.8
##  young     7.8 4.2

round(ct$p.value, 4)      # Yates-corrected

## [1] 0.0098

ft <- fisher.test(tab)    # exact test
round(ft$p.value, 4)

## [1] 0.0044
```

Reading the categorical analysis

```
tab_g <- with(data4, table(Gender, Sugar))  
ft_g <- fisher.test(tab_g)
```

- **Age:** Fisher's exact test gives $p = 0.0044$, well below 0.05. Sugar is associated with age group: 6 of 8 old patients against 1 of 12 young ones.
- **Gender:** the same test gives $p = 0.374$: no evidence of an association in these data.
- The chi-square test agrees for age ($p = 0.0098$), but its p -value is only approximate with so few patients.

What a p -value says The probability, if H_0 were true, of a table at least as far from independence as the one observed. It is not the probability that H_0 is true.

Caution Association is not causation, and 20 hypothetical patients cannot represent a population.

- 1 Getting started
- 2 Vectors and data
- 3 Graphics
- 4 Categorical data
- 5 Comparing groups**
- 6 Correlation and regression

Two independent batches: the question

Scores of five students in each of two batches of the same course. Do the two batches have the same mean score?

$$H_0 : \mu_A = \mu_B \quad \text{vs} \quad H_1 : \mu_A \neq \mu_B$$

$$\text{Welch: } t = \frac{\bar{x}_A - \bar{x}_B}{\sqrt{s_A^2/n_A + s_B^2/n_B}}$$

Welch's test does not assume equal variances; its degrees of freedom are estimated from the data, so they need not be a whole number.

```
A1 <- c(20, 20.5, 13, 16, 11) # batch A
B1 <- c(40, 42, 38, 56, 57)   # batch B
c(mean(A1), mean(B1))

## [1] 16.1 46.6

c(sd(A1), sd(B1))

## [1] 4.189 9.154
```

First look Batch B has the higher mean and the larger spread. Is the gap larger than chance variation between students would produce?

Welch's two-sample t -test in R

```
t.test(A1, B1)           # Welch's test is the default

##
##  Welch Two Sample t-test
##
## data:  A1 and B1
## t = -6.8, df = 5.6, p-value = 0.0007
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##   -41.71 -19.29
## sample estimates:
## mean of x mean of y
##      16.1      46.6
```

The output gives the statistic, its degrees of freedom and p -value, a 95% confidence interval for $\mu_A - \mu_B$ and the two sample means.

The pooled t -test, and what `t.test()` returns

Assuming $\sigma_A^2 = \sigma_B^2$:

$$s_p^2 = \frac{(n_A - 1)s_A^2 + (n_B - 1)s_B^2}{n_A + n_B - 2}$$

$$t = \frac{\bar{x}_A - \bar{x}_B}{s_p \sqrt{1/n_A + 1/n_B}} \sim t_{n_A + n_B - 2}$$

The result is a list: `names(fit.t)` lists its ten parts, and `$` picks one out.

```
fit.t <- t.test(A1, B1, var.equal = TRUE)
fit.t$statistic; fit.t$parameter

##          t
## -6.774
## df
##  8

fit.t$p.value

## [1] 0.0001415

fit.t$conf.int

## [1] -40.88 -20.12
## attr(,"conf.level")
## [1] 0.95
```

Reading the two-sample tests

- Mean scores: batch A 16.1, batch B 46.6; difference -30.5 points
- Welch: $t = -6.77$, $p = 0.0007$; 95% CI for $\mu_A - \mu_B$: -41.7 to -19.3
- Pooled: $t = -6.77$ on 8 df, $p = 0.0001$, CI -40.9 to -20.1
- Both reject H_0 : batch B scored higher on average

Which one? Welch is R's default and is safe whether or not the variances are equal (here $s_A = 4.2$, $s_B = 9.2$).

The interval We are 95% confident that batch B's mean exceeds batch A's by between 19 and 42 points; the interval excludes 0.

Paired t -test: if the same five students sat both tests

$$d_i = x_{Ai} - x_{Bi}, \quad H_0 : \mu_d = 0$$

$$t = \frac{\bar{d}}{s_d/\sqrt{n}} \sim t_{n-1}$$

Reading it Mean difference -30.5 points, $t = -5.81$ on 4 df, $p = 0.0044$. Pair only when the design pairs the data.

```
t.test(A1, B1, paired = TRUE)

##
## Paired t-test
##
## data: A1 and B1
## t = -5.8, df = 4, p-value = 0.004
## alternative hypothesis: true mean difference is not equal to 0
## 95 percent confidence interval:
## -45.09 -15.91
## sample estimates:
## mean difference
## -30.5
```

The formula interface and a data frame

```
score <- c(A1, B1)
group <- gl(2, 5, labels = c("batchA",
                             "batchB"))
data6 <- data.frame(score, group)
head(data6, 3)

##   score group
## 1  20.0 batchA
## 2  20.5 batchA
## 3  13.0 batchA

t.test(score ~ group, data = data6)$p.value

## [1] 0.0006718
```

- A **data frame** is a list of equal-length columns: one row per student
- `score ~ group` reads “score explained by group”
- Same Welch test, same p -value as before

Why bother? The same formula style is used by `aov()`, `lm()`, `boxplot()` and many more.

Three batches: one-way analysis of variance

$$y_{ij} = \mu + \tau_i + \varepsilon_{ij}, \quad \varepsilon_{ij} \sim N(0, \sigma^2)$$

$$H_0 : \tau_A = \tau_B = \tau_C = 0$$

$$F = \frac{MS_{\text{between}}}{MS_{\text{within}}} \sim F_{k-1, n-k}$$

Comparing three means with three separate t -tests would inflate the chance of a false positive; ANOVA tests them together.

```
C1 <- c(10, 15, 12, 10, 3)      # batch C
data7 <- data.frame(
  score = c(A1, B1, C1),
  group = gl(3, 5, labels = c("batchA",
                              "batchB", "batchC"))
)
M2 <- aov(score ~ group, data = data7)
print(summary(M2), signif.stars = FALSE)
```

##	Df	Sum Sq	Mean Sq	F value	Pr(>F)
## group	2	3845	1923	47.7	0.0000019
## Residuals	12	483	40		

Group means and Tukey's multiple comparisons

```
model.tables(M2, type = "means")
```

```
## Tables of means
## Grand mean
##
## 24.23
##
## group
## group
## batchA batchB batchC
## 16.1 46.6 10.0
```



```
TukeyHSD(M2, which = "group")
```

```
## Tukey multiple comparisons of means
## 95% family-wise confidence level
##
## Fit: aov(formula = score ~ group, data = data7)
##
## $group
## diff lwr upr p adj
## batchB-batchA 30.5 19.79 41.209 0.0000
## batchC-batchA -6.1 -16.81 4.609 0.3167
## batchC-batchB -36.6 -47.31 -25.891 0.0000
```

Tukey's HSD compares every pair of means while keeping the family-wise error rate at 5%.

Seeing the ANOVA

```
par(mfrow = c(1, 2), mar = c(4, 7, 2, 1), cex.main = 0.9)
boxplot(score ~ group, data = data7, col = col_main, las = 1, horizontal = TRUE,
        ylab = "", xlab = "Score", main = "Scores by batch")
plot(TukeyHSD(M2, which = "group"), las = 1)
```



Right: 95% family-wise intervals for each difference. An interval that crosses 0 (C-A) means those two batches cannot be distinguished.

Reading the ANOVA

- $F = 47.7$ on 2 and 12 df, $p < 0.001$: the three batch means are not all equal
- Tukey: B is higher than A by 30.5 and higher than C by 36.6 points (both adjusted $p < 0.001$)
- A and C differ by 6.1 points, adjusted $p = 0.32$: no evidence of a difference

In one sentence Batch B outperformed both other batches; batches A and C performed alike.

Assumptions Independent observations, roughly normal errors and similar spreads in the groups. With five students per batch, look at the box plots before trusting the test.

- 1 Getting started
- 2 Vectors and data
- 3 Graphics
- 4 Categorical data
- 5 Comparing groups
- 6 Correlation and regression**

Correlation between marks

```
data8 <- read.table("data/data8.txt",  
                    header = TRUE)
```

```
data8
```

```
##   batchA batchB batchC  
## 1   20.0    40    10  
## 2   20.5    42    15  
## 3   13.0    56    12  
## 4   16.0    38    10  
## 5   11.0    57     3
```

```
round(cor(data8), 3)
```

```
##           batchA batchB batchC  
## batchA   1.000 -0.830  0.696  
## batchB  -0.830  1.000 -0.476  
## batchC   0.696 -0.476  1.000
```

Pearson's correlation coefficient

$$r = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2 \sum (y_i - \bar{y})^2}}$$
$$-1 \leq r \leq 1$$

Pairs needed Correlation needs two measurements on the *same* units. Read each row as one student's marks in three papers (illustrative data).

Testing a correlation

```
ct8 <- cor.test(data8$batchA, data8$batchB)
ct8$estimate; ct8$p.value

##      cor
## -0.8299
## [1] 0.08206

ct8$conf.int[1:2]      # 95% CI for rho

## [1] -0.9884  0.1956

cor(data8$batchA, data8$batchB,
     method = "spearman")

## [1] -0.6
```

$$H_0 : \rho = 0, \quad t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}} \sim t_{n-2}$$

Reading it $r = -0.83$ looks strong, yet $p = 0.082$ and the 95% CI for ρ runs from -0.99 to 0.2 . With $n = 5$ we cannot rule out zero correlation. Spearman's rank correlation is -0.6 .

Fitting a simple linear regression

$$y_i = \beta_0 + \beta_1 x_i + \varepsilon_i, \quad \varepsilon_i \sim N(0, \sigma^2)$$

Least squares: choose $\hat{\beta}_0, \hat{\beta}_1$ to minimise $\sum (y_i - \hat{y}_i)^2$


`lm()` fits the model; `summary()` gives the full table and `display()` from `arm` a compact one.

```
M1 <- lm(batchA ~ batchB, data = data8)
display(M1, detail = TRUE)

## lm(formula = batchA ~ batchB, data = data8)
##               coef.est coef.se t value Pr(>|t|)
## (Intercept)  33.80      6.98   4.85    0.02
## batchB       -0.38      0.15  -2.58    0.08
## ---
## n = 5, k = 2
## residual sd = 2.70, R-Squared = 0.69
```

Reading the regression

```
par(mar = c(4, 4, 1, 1))  
plot(batchA ~ batchB, data = data8, pch = 16,  
      col = col_main, cex = 1.3)  
abline(M1, col = col_alt, lwd = 2)
```



- Fitted line: $\hat{A} = 33.8 - 0.38 \times B$
- Each extra mark in paper B goes with 0.38 fewer marks in paper A, on average
- $R^2 = 0.69$, but the slope has $p = 0.082$ and $r = -0.83$ is not significant at 5%

Honest conclusion Five points and 3 residual df cannot establish a relationship. Use this example to learn the commands, not to draw conclusions.

Key take-aways

- 1 **Work in a script**, inside an RStudio project, with relative paths.
- 2 **Vectors are the building blocks**: functions act on every element at once.
- 3 **Look at the data first**: tables, proportions and plots before tests.
- 4 **Check a test's conditions**: small expected counts call for Fisher's exact test.
- 5 **Let the design choose the test**: independent groups or paired measurements.
- 6 **Report effect sizes with intervals**, not only p -values, and remember that tiny samples give weak evidence.

- Dalgaard, P. (2008). *Introductory Statistics with R*, 2nd Ed., Springer, New York.
- Gelman, A., and Hill, J. (2007). *Data Analysis Using Regression and Multilevel/Hierarchical Models*, Cambridge University Press, Cambridge.
- Ihaka, R., and Gentleman, R. (1996). R: A language for data analysis and graphics. *Journal of Computational and Graphical Statistics*, 5(3), 299–314.
- Jones, O., Harden, S., and Crawley, M. J. (2022). *The R Book*, 3rd Ed., Wiley, Hoboken.
- Verzani, J. (2014). *Using R for Introductory Statistics*, 2nd Ed., CRC Press, Boca Raton.
- R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna. <https://www.R-project.org>

Thank you

Questions and discussion